

Piedmont Stone Center Protocols

Non-Steroidal Anti-inflammatory Drugs (NSAIDs)

Discontinue 48 hours prior to Lithotripsy

A	D	I	N	S
Actiprofen	Daypro	Ibuprofen	Naxen	Salgesic
Actron	Diclofenac	IBU Tabs	NeoProfen	Salonpas
Advil	Diclofenac Gel	Indameth	Novo-Difenac	Salsalate
Aleve	Dicloflex	Indocid	Novo-Naprox	Salsitab
Amigesic	Difenac	Indocin	Novo-Sundac	Sampson-8 Select
Anaflex	Diflunisal	Indomethiacin	Novomethacin	Sandoz Diclofenac
Anaprox	Dimetapp Sinus	K/L	Novopirocam	Serophene
Anaprox DS	Disalcid	Ketoprofen	Novoprofen	Sine-Aid IB
Anodynos	Doan's Pills	Ketoprofen ER	Nu-Diclo	Solaraze
Ansaid	Dolobid	<u>Ketorolac – Stop for 24 hours</u>	Nu-Ibuprofen	Sprix Nasal Spray
Apo-Dilo	Dristan Sinus	Lodine, Lodine XL	Nu-Indo	Sulindac
Apo-Ibuprofen	Duexis	M	Nu-Naprox	Synflex
Apo-Indomethiacin	Durasol		Nuprin	Synflex DS
Apo-Keto	Dyloject	Magan	O	T
Apo-Meloxicam	E	Mathritic	Orafen	Tab-profen
Apo-Nabumetone	ElixSure-IB	Meclofenamate	Oruvail	Tolectin
Apo-Naproxen	Equaline All Day	Meclomen	Orudis	Tolmetin
Apo-Napro-NA	Relief	Medipren	Oxaprozin	Top Care All Day
Apo-Oxaprozin	Excedrine IB	Mefenamic Acid	P	Relief
Apo-Piroxicam	Etodolac	Meloxicam	Pamprin	<u>Toradol – Stop for 24 hours</u>
Aposulin	F	Menadol	Pennsaid	Treximet
Arthrotec	Feldene	Midol	Phenylbutazone	V
Azolid	Fenoprophen	Midol Extended Relief	Pirocam	Valdecocib
	Flector	Mobic	Piroxicam	Vicoprofen
	Flurbiprophen	Mobidin	PMS-Pirocam	Vimovo
	Froben	Mobicox	Ponstel	Voltaren
	G	Mono-Gesic	Profen	Voltaren Gel
	Gen-Meloxicam	Motrin IB	R	W
	Gen-Nabumetone	Motrin	Rapide	Walprofen
	Gen-Naproxen	N	Relafen	Walproxen
	Gen-Piroxicam	Nabumetone	Reprexain	X
	Genpril	Nalfon	Rhodis	Z
	Good Sense Pain	Naprelan	Rhovail	
	Relief	Naprosyn	Rufen	
	H	Naprosyn EC	S	
	Haltran	Naproxen	Salflex	

Anticoagulant and Antiplatelet Lab Requirements

Consult with patient's cardiologist before stopping any of the medications below

Arixtra (Fondaparinux)	Discontinue 3 days (72 hours) prior to treatment. Need/draw CBC, platelet labs. Check creatinine for patients 75 and older. Labs must be WNL.
Brilinta (Ticagrelor)	Discontinue for 5 days (120 hours) prior to treatment.
Coumadin (Warfarin, Warfilone)	Discontinue, treat when PT (prothrombin) or INR is normal. INR must be 1.4 or less. If patient is off Coumadin for 3 weeks, no labs necessary.
Desirudin (Iprivask)	Discontinue 1 day (24 hours) and check PTT. PTT should be normal.
Effient (Prasugrel)	Discontinue for 7 days (168 hours) prior to treatment. <i>Check with prescribing physician first!</i>
Eliquis (Apixiban)	Discontinue for 2 days (48 hours) prior to treatment. No labs required.
Elmiron	Discontinue for 3 days (72 hours) prior to treatment.
Fragmin (Dalteparin)	Discontinue 1 day (24 hours) prior to treatment.
Heparin	Draw PTT if last dose within 6 hours of treatment.
Innohep (Tinzaparin)	Discontinue 1 day (24 hours) prior to treatment.
Integrillin (Eptifibatide)	Discontinue 12 hours prior to treatment. This is an IV medication, consult with cardiologist regarding stopping the medication.
Jantoven (same as Coumadin)	Discontinue, treat when PT (prothrombin) or INR is normal. INR must be 1.4 or less. If patient is off Jantoven for 3 weeks, no labs necessary.
Lovenox (Enoxaparin)	Discontinue for 24 hours prior to treatment.
Persantine (Dipyridamole)	Discontinue for 3 days (72 hours) prior to treatment.
Plavix (Clopidogrel)	Discontinue for 5 days (120 hours) prior to treatment.
Pletol (Cilostazol)	Discontinue for 4 days (96 hours) prior to treatment.
Pradaxa	Discontinue for 5 days (120 hours) prior to treatment. If less than 5 days, draw PTT. PTT should be WNL.
Ticlid (Ticlopidine HCL)	Discontinue for 14 days (336 hours) prior to treatment. Need CBC, platelet labs 1-2 days prior to treatment.
Trental (Pentoxifylline)	Discontinue for 3 days (72 hours) prior to treatment.
Tyvaso (Treprostinil, Remodulin)	Discontinue for 2 days (24 hours) prior to treatment.
Vorapaxar (Zontivity)	<i>Check with physician first!</i>
Xarelto (Rivaroxaban)	Discontinue for 3 days (72 hours) prior to treatment. If less than 72 hours, need/draw PTT. PTT should be WNL.

