

Piedmont Stone Center Protocols
Aspirin Products / Naltrexone/Suboxone
Discontinue 72 hours prior to Lithotripsy

<p>A</p> <p>Acetasalicylic Acid Acuprin Aggrenox Alka Seltzer Alka Seltzer Plus Alor Amigesic Anacin Apo-ASA Apo-Asen Argesic Arthrinol Arthritis Arthritis Pain Formula Arthritis Strength Bufferin Arthropan Liquid Arthrotec Artria AR ASA ASA Enscales Ascriptin Ascriptin A/D Ascriptin with Codeine Asperbuf Aspercreme Aspergum Aspir-Low Aspirin Aspir-trin Aspiatab Astrin Atromid Axotal</p> <p>B</p> <p>Bayer Aspirin Bayer Children's Aspirin Bayer Children's Cold Tablets Bayer Muscle & Joint Cream Bayer Time Release Tablets BC Tablets and Powders Bismatrol Bismed Bismylate Bismuth Subsalsicylate Buffetts II Tablets Buff -A Comp Tablets Bufferin with or w/o Codeine Butalbital-ASA (Axotal)</p> <p>C</p> <p>Cama Arthritis Pain Reliever Carisprodol-ASA Choline Magnesium Trisalsicylate Choline Salsicylate Claripex CMT Congesprin</p>	<p>C</p> <p>Cope Tablets Coryphen Cosprin Tablets CVS Arthricream Cream Rub CVS Muscle Rub CVS Muscle Rub Ultra Strength</p> <p>D</p> <p>Damason-P Darvon Compound Darvon Compound 65 Darvon Compound with ASA Digesic Disalcid</p> <p>E</p> <p>Easprin Ecotrin Efficin Emagrin Empirin with Codeine Emprazil Encaprin Endodan Entrophen Equagesic Esaprin Excedrine Excedrine Migraine</p> <p>F</p> <p>Fiorinal Fiorinal with Codeine Fiorial with Codeine Flexall Plus Froben</p> <p>G</p> <p>Gelprin Gemnisyn Goodys</p> <p>H</p> <p>Halfprin Helidac</p> <p>K</p> <p>Kaopectate Kao-Tin</p> <p>L</p> <p>Lanorinal Lortab ASA</p> <p>M</p> <p>Magan Magnaprin Magsal Marnal Maximum Bayer Aspirin Measurin Mesasal Methocarbonal with Aspirin Micranin</p>	<p>M</p> <p>Mobidin Mobigesic Momentum Monogesic Myogesic</p> <p>N</p> <p>Neoccybate Norgesic Norgesic Forte Norwich Novasen Novo-Fibrate</p> <p>O</p> <p>Orphengesic Orphengesic Forte Oxycodan</p> <p>P</p> <p>Pabalate SF P-A-C Analgesic Panasal 5/500 Pepto-Bismol Peptic-Relief Percodan Percodan-Demi Persantine Persistin Phenaphen with Codeine Pink Bismuth PMS-ASA Praviguard PAC Propoxyphene Compound 65</p> <p>R</p> <p>Rexolate Robaxisol Roxiprin</p> <p>S</p> <p>Saleto Tablets SK 65 Compound Sloprin Sodium Thiosalsicylate Soma Compound St. Josephs Aspirin Sportscreme Stanback Supasa Synalgos-DC</p> <p>T</p> <p>Talwin Tecnal Teva- Oxycodan Triaminicin Tricosal Trigesic Trilisate</p> <p>V</p> <p>Vanquish Vazalore Z- Zorprin</p>
--	---	---

Naltrexone (Vivitrol, Revia) - Advised to discontinue for 72 hours – check with your prescribing MD before stopping. If you cannot stop your Naltrexone, please let your urologist know ASAP.

Suboxone/Buprenorphine – Advise to discontinue for 72 hours – check with your prescribing MD before stopping. If you cannot stop your Suboxone, please let your urologist know ASAP.

Herbal Medicines –Advised to discontinue for 1 week prior to lithotripsy.