

# PIEDMONT STONE CENTER

## Non-Steroidal Anti-inflammatory Drugs (NSAIDs)

### Discontinue 48 hours prior to Lithotripsy

A	D	I	N	S
Actiprofen	Daypro	Ibuprofen	Naxen	Salgesic
Actron	Diclofenac – <b>pill only</b>	IBU Tabs	NeoProfen	Salonpas Arthritis
Advil	Dicloflex – <b>pill only</b>	Indameth	Novo-Difenac	<b>(with Methyl Salicylate)</b>
Aleve	Difenac – <b>pill only</b>	Indocid	Novo-Naprox	Salsalate
Amigesic	Diflunisal	Indocin	Novo-Sundac	Salsitab
Anaflex	Dimetapp Sinus	Indomethiacin	Novomethacin	Sampson-8 Select
Anaprox	Disalcid	<b>K/L</b>	Novopirocam	Sandoz Diclofenac
Anaprox DS	Doan's Pills	Ketoprofen	Novoprofen	Serophene
Anodynol	Dolobid	Ketoprofen ER	Nu-Diclo	Sine-Aid IB
Ansaid	Dristan Sinus	<b><u>Ketorolac – Stop for 24 hours</u></b>	Nu-Ibuprofen	Solaraze
Apo-Dilo	Duexis	<b><u>24 hours</u></b>	Nu-Indo	Sprix Nasal Spray
Apo-Ibuprofen	Durasol	Lodine, Lodine XL	Nu-Naprox	Sulindac
Apo-Indomethiacin	Dyloject	<b>M</b>	Nuprin	Synflex
Apo-Keto	<b>E</b>	Magan	<b>O</b>	Synflex DS
Apo-Meloxicam	ElixSure-IB	Mathritic	Orafen	<b>T</b>
Apo-Nabumetone	Equaline All Day	Meclofenamate	Oruvail	Tab-profen
Apo-Naproxen	Relief	Meclomen	Orudis	Tolectin
Apo-Napro-NA	Excedrine IB	Medipren	Oxaprozin	Tolmetin
Apo-Oxaprozin	Etodolac	Mefenamic Acid	<b>P</b>	Top Care All Day
Apo-Piroxicam	<b>F</b>	Meloxicam	Pamprin	Relief
Aposulin	Feldene	Menadol	Pennsaid	<b><u>Toradol – Stop for 24 hours</u></b>
Arthrotec	Fenoprophen	Midol	Phenylbutazone	<b>hours</b>
Azolid	Flector	Midol Extended Relief	Pirocam	Treximet
<b>B</b>	Flurbiprophen	Mobic	Piroxicam	<b>V</b>
Butazolidin	Froben	Mobidin	PMS-Pirocam	Valdecoxib
<b>C</b>	<b>G</b>	Mobicox	Ponstel	Vicoprofen
Caldolor	Gen-Meloxicam	Mono-Gesic	Profen	Vimovo
Cambia	Gen-Nabumetone	Motrin IB	<b>R</b>	Voltaren – <b>pill only</b>
Cataflam	Gen-Naproxen	Motrin	Rapide	<b>W</b>
Combunox	Gen-Piroxicam	<b>N</b>	Relafen	Walprofen
Clinoril	Genpril	Nabumetone	Reprexain	Walproxen
Clomiphene	Good Sense Pain Relief	Nalfon	Rhodus	<b>X</b>
<b>H</b>	Haltran	Naprelan	Rhovail	<b>Z</b>
		Naprosyn	Rufen	Zipsor
		Naprosyn EC	Salflex	Zorvolex
		Naproxen		

### Anticoagulant and Antiplatelet Lab Requirements

*Consult with patient's cardiologist before stopping any of the medications below*

Arixtra (Fondaparinux)	Discontinue 3 days (72 hours) prior to treatment. Need/draw CBC, platelet labs. Check creatinine for patients 75 and older. Labs must be WNL.
Brilinta (Ticagrelor)	Discontinue for 5 days (120 hours) prior to treatment.
Coumadin (Warfarin, Warfilone)	Discontinue for 5 days. Treat when PT (prothrombin) or INR is normal. INR must be 1.4 or less. If patient is off Coumadin for 3 weeks, no labs necessary.
Desirudin (Iprivask)	Discontinue 1 day (24 hours) and check PTT. PTT should be normal.
Effient (Prasugrel)	Discontinue for 7 days (168 hours) prior to treatment. <b>Check with prescribing physician first!</b>
Eliquis (Apixiban)	Discontinue for 2 days (48 hours) prior to treatment. No labs required.
Elmiron	Discontinue for 3 days (72 hours) prior to treatment.
Fragmin (Dalteparin)	Discontinue 1 day (24 hours) prior to treatment.
Heparin	Draw PTT if last dose within 6 hours of treatment.
Innohep (Tinzaparin)	Discontinue 1 day (24 hours) prior to treatment.
Integrillin (Eptifibatide)	Discontinue 12 hours prior to treatment. This is an IV medication, consult with cardiologist regarding stopping the medication.
Jantoven (same as Coumadin)	Discontinue, treat when PT (prothrombin) or INR is normal. INR must be 1.4 or less. If patient is off Jantoven for 3 weeks, no labs necessary.
Lovenox (Enoxaparin)	Discontinue for 24 hours prior to treatment.
Persantine (Dipyridamole)	Discontinue for 3 days (72 hours) prior to treatment.
Plavix (Clopidogrel)	Discontinue for 5 days (120 hours) prior to treatment.
Pletol (Cilostazol)	Discontinue for 4 days (96 hours) prior to treatment.
Pradaxa	Discontinue for 5 days (120 hours) prior to treatment. If less than 5 days, draw PTT. PTT should be WNL.
Ticlid (Ticlopidine HCL)	Discontinue for 14 days (336 hours) prior to treatment. Need CBC, platelet labs 1-2 days prior to treatment.
Trental (Pentoxifylline)	Discontinue for 3 days (72 hours) prior to treatment.
Tyvaso (Trepstinil, Remodulin)	Discontinue for 2 days (24 hours) prior to treatment.
Vorapaxar (Zontivity)	<b>Check with physician first!</b>
Xarelto (Rivaroxaban)	Discontinue for 3 days (72 hours) prior to treatment. If less than 72 hours, need/draw PTT. PTT should be WNL.